

PEDIATRIC INFECTIOUS
DISEASE SOCIETY OF THE
PHILIPPINES

PIDSP JOURNAL

Vol.13 No.2 July-Dec 2012

Dengue: A Growing Global Health Threat Usa Thisyakorn M.D., Chule Thisyakorn, M.D. Department of Pediatrics,
Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand2
Original Articles A Case Control Study of the Demographic Characteristics, Risk Factors Clinical Findings, Treatment and Outcome among Children 18 Years and Below Who are Confirmed to have Influenza A H1N1/09 Virus Romina D. Gerolaga, M.D., Robert Dennis Garcia, M.D. Makati Medical Center
The Clinical Outcome and Antibiotic Sensitivity Pattern of Enterobacter Spp. Culture Positive Neonates Admitted at Cebu Doctors' University Hospital – Neonatal Intensive Care Unit (2005-2008) Lalaine Amor H. Maderal, M.D.*, Barbra Charina V. Cavan, M.D.* Cebu Doctors' University Hospital
Haemophilus Influenzae Type B Conjugate Vaccine (HiBCV) and Heptavalent Pneumococcal Conjugate Vaccine (PCV7) Immunization Status Of Patients 5 Years and Below Hospitalized For Pneumonia Lou Ver Leigh A. Manzon, M.D., Robert Dennis J. Garcia, M.D., Sally Victoria B. King, M.D Department of Pediatrics, Makati Medical Center30
Interim Advice Post Disaster Interim Advice on the Prevention of Leptospirosis in Children
Outbreak of Serratia Marcescens in the Newborn Care Unit in a Local Tertiary Hospital Sandra Joyce Pena, MD, Xenia Cathrine Fabay, MD Baguio General Hospital and Medical Center
The Use of Meropenem among Neonates: A One-Year Retrospective Study in the Nursery of a Local Tertiary Hospital Ferdinand P. Ganggangan, MD, Xenia Cathrine Fabay, MD Baguio General Hospital and Medical Center
PIDSP Newsletter PIDSP AT THE APAME CONGRESS 201254 PIDSP Joins the PPS Total Family Healthcare Expo56

Vol.13 No.2 July-Dec 2012

POST DISASTER INTERIM ADVICE ON THE PREVENTION OF LEPTOSPIROSIS IN CHILDREN

by the

Pediatric Infectious Disease Society of the Philippines, Inc.

August 10, 2012

The purpose of this advice is to provide guidelines for physicians and parents on the prevention of leptospirosis in the pediatric age group.

Leptospirosis is an infectious disease caused by pathogenic bacteria called leptospires. The main carriers of leptospires are animals such as rats, mice, dogs, cats and livestock. Human leptospiral infections can occur when mucus membranes and skin are contaminated by the urine of infected animals, or upon ingestion of contaminated food and water, all of which may occur when an individual is exposed to flood waters. Thus the occurrence of freshwater flooding after typhoons or increased rainfall can lead to an increased risk of leptospirosis.

Initial symptoms of leptospirosis may range from mild to severe, or may even be fatal. Manifestations of the disease include: (1) a mild influenza-like illness; (2) Weil's syndrome, characterized by jaundice, renal failure, hemorrhage, and arrhythmias; (3) meningitis/meningoencephalitis; or (4) pulmonary hemorrhage with respiratory failure. Case fatality rates range from 12-14% in the Philippines. In studies in India, case fatality rates were lower in the pediatric age group compared to adults.

Prevention of leptospirosis includes the following:

- 1. Parents should instruct children not to wade or swim in flood waters.
- 2. If exposure to flood waters is unavoidable, protective gear such as boots, goggles, overalls, and rubber gloves should be used.
- 3. All food and drinking water should be protected against contamination. Fresh vegetables and fruit should be washed in previously boiled or clean water and then cooked or peeled.
- 4. Boil drinking water for at least 10-15 minutes. Physical filtration through ceramic or charcoal filters is not adequate for leptospirosis.
- 5. Food should be protected against rodent attack or contamination.
- 6. If children are exposed to flood waters, antibiotic prophylaxis may decrease occurrence of clinical disease and mortality. Prophylactic antibiotics should be given under the supervision of a physician, who can give advice regarding effects, precautions and contraindications for these medications.

The following antibiotics are recommended for children exposed to flood waters: Drug of Choice

	Dose (oral)	Comments
Doxycycline	4 mg/kg single dose, max dose: 200mg	Proven efficacy for preventing clinical disease Adverse effects are similar to other tetracyclines; in children below 8 years of age, doxycycline is unlikely to cause dental staining at the dose and duration recommended to treat serious infections Avoid milk, dairy products, iron and antacids 1 hour before and 2 hours after administration; may be given with food to avoid stomach upset

ALTERNATIVE DRUGS

	Dose (oral)	Comments
Azithromycin	10 mg/kg single dose, max dose: 500 mg	Efficacy for prevention of leptospirosis was seen in in vitro and animal models
Amoxicillin	50 mg/kg/day q 6 hrs. for 3-5days Max dose: 500 mg q 6 hrs.	No clinical trial for prevention of leptospirosis, but amoxicillin is a known alternative for the treatment of disease Dose is for 3-5 days due to the very short half-life

If children are exposed for more than 7 days, the dose should be repeated after 1 week. Prophylaxis is not 100% effective. Prevention of exposure is most prudent. Monitor all those exposed for the occurrence of symptoms of leptospirosis. The early signs of infection occur between 4 and 10 days after exposure, and usually begin with a fever and an influenza-like illness. Headaches, sensitivity to light, muscle and joint pains, vomiting and fatigue are also common. Seek medical attention once these symptoms are noted.

References:

- 1. The Leptospirosis Task Force (PSMID, PSN, PCP) Philippine Clinical Practice Guidelines on the Diagnosis, Management and Prevention of Leptospirosis 2010
- 2. Amilasan A, Ujiie M et al, Outbreak of Leptospirosis after Flood, the Philippines, 2009. Emerging Infectious Diseases, 2012; 18(1): 91-94.
- 3. Leptospirosis in the Philippines, updated October 2009. http://www.gideononline.com/tag/leptospirosis/ accessed on August 8, 2012
- 4. Sehgal SC, Sugunan AP, Murhekar MV, Sharma S, Vijayachari P. Randomized controlled trial of doxycycline prophylaxis against leptospirosis in an endemic area. International Journal of Antimicrobial Agents 2000; 13 (4): 249-255.
- 5. Illangasekera VL, Kularatne SA, Kumarasiri PV, Pussepitiya D, Premaratne MD. Is oral penicillin an effective chemoprophylaxis against leptospirosis? A placebo controlled field study in the Kandy District, Sri Lanka. Southeast Asian J Trop Med Public Health 2008; 39(5):882-4
- 6. Antony J, Celine TM, Chacko M. Case fatality rate of leptospirosis in a tertiary care hospital in Kerala, India. Ann Trop Med Public Health [serial];5:236-239
 - Available from: http://www.atmph.org/text.asp?2012/5/3/236/98626
 - Accessed on Aug 8, 2012
- 7. Lopes AA, Costa E, Costa YA, et al. Comparative Study Of The In-Hospital Case-Fatality Rate Of Leptospirosis Between Pediatric And Adult Patients Of Different Age Groups. Rev. Inst. Med. trop. S. Paulo 2004: 46(1):19-24
- 8. Ghouse M, Maulana AB, Mohamed Ali MG, Sarasa VD. A two-year study of the efficacy of treatment of leptospirosis in humans. Indian J Med Microbiol 2006; 24:345-6
- 9. Griffith ME, Hospenthal DR, Murray CK. Antimicrobial therapy of leptospirosis. Curr Opin Infect Dis. 2006 Dec; 19(6):533-7.
- 10. Kriangsak Phimda, Siriwan Hoontrakul,et al. Doxycycline versus Azithromycin for Treatment of Leptospirosis and Scrub Typhus. *Antimicrob Agents Chemother*. 2007, 51(9):3259
- 11. Hospenthal D, Murray CK. In Vitro Susceptibilities of Seven *Leptospira* Species To Traditional And Newer Antibiotics. Antimicrobial Agents And Chemotherapy 2003; 47 (8): 2646–2648.
- 12. Moon JE, Rivard RG. Effect of timing and duration of azithromycin therapy of leptospirosis in a hamster model. Journal of Antimicrobial Chemotherapy (2007) 59, 148–151
- 13. Takafuji ET, Kirkpatrick JW. An Efficacy Trial of Doxycycline Chemoprophylaxis against Leptospirosis. N Engl J Med 1984; 310:497-500
- 14. Milind S, Tullu, Sunil Karande. Leptospirosis in children: A review for family physicians. Indian Journal of Medical Sciences, 2009; 63 (8): 368-378.
- 15. Everett ED. Treatment and prevention of leptospirosis. Official Reprint from UptoDate.com. http://www.uptodate.com/contents/treatment-and-prevention-of-leptospirosis?view=print[8/9/2012 1:39:27 PM]